

Front panel

Back panel

PADCEV™
(enfortumab vedotin)

PATIENT CARD



GLUE SPOT
TEXT FREE

- Carry this card with you **at all times**, especially when you travel or when you see another doctor.
- Please ensure you show this card to any doctor, pharmacist or nurse for any medical treatment or at any visits to the hospital or clinic.
- Please contact your doctor **immediately**, if you develop any side effects, in particular those listed on this card.

**IMPORTANT SAFETY
INFORMATION FOR
PATIENTS**

Padcev may cause serious side effects, including severe skin reactions (Stevens-Johnson syndrome (SJS), Toxic Epidermal Necrolysis (TEN) and other severe rashes such as symmetrical drug related intertriginous and flexural exanthema).

Talk to your doctor, pharmacist or nurse **immediately** if you have any of the following symptoms:

- rash or itching that continues to get worse or comes back after treatment,
- skin blistering or peeling,
- painful sores or ulcers in mouth or nose, throat, or genital area,
- fever or flu like symptoms,
- or swollen lymph nodes.

These may be signs of a severe skin reaction that can happen while receiving this medicine, particularly during the first few weeks of treatment. If it occurs, your doctor will monitor you and may

give you medicine to treat your skin condition. She or he may pause or stop treatment if your skin reaction worsens. If you have any further questions about your treatment, please contact your doctor.

**IMPORTANT
INFORMATION
FOR HEALTHCARE
PROVIDERS**

- This patient is being treated with Padcev (enfortumab vedotin), which can cause severe skin reactions, including SJS and TEN

(predominantly during the first cycle of treatment).

- Symptoms include rash or itching that continues to get worse or comes back after treatment, skin blistering or peeling, painful sores or ulcers in mouth or nose, throat, or genital area, fever or flu-like symptoms or swollen lymph nodes.
- Fever or flu-like symptoms may be the first sign of a skin reaction. Patients should be monitored starting with the first cycle and throughout treatment for skin reactions.

Topical corticosteroids/ antihistamines can be considered for mild to moderate skin reactions.

- If SJS or TEN is suspected or if bullous lesions occur, **immediately** withhold treatment and refer for specialised care; histologic confirmation is critical to early recognition, as diagnosis and intervention can improve prognosis.
- If SJS or TEN, Grade 4 or recurrent Grade 3 skin reactions occur, permanently discontinue treatment.

- Withhold treatment for Grade 2 with fever, worsening Grade 2 or Grade 3 skin reactions until Grade ≤ 1 and resume at the same dose level or consider dose reduction by one dose level; consider referral to specialised care.

Please contact the patient's Oncologist for more information and consult the Product Information for enfortumab vedotin available at www.medicines.org.uk.

My name:

My contact number:

Emergency contact:

**Emergency contact
number:**

**Name of Oncologist/
Oncology Nurse:**

Contact number:

**After-hours contact
number:**

Name of my Hospital:

**My Hospital contact
number:**

PADCEV start date:

This card was last revised
in 04/2022.